

CASE REPORT

Fat Embolism Syndrome Complicated by Acute Pulmonary Thromboembolism after Bilateral femoral shaft Fractures: Two Nightmares in the Same Patient

Aamir Shafi, JawharUl Islam, Wani Abdul Ahad, Aariba Zahoor

Abstract

Fat embolism syndrome is an uncommon but fatal complication usually in orthopedic trauma especially after a long bone fracture. A high level of suspicion should be kept in mind when a patient of long bone fracture develops hypoxia, confusion or rash in the absence of infection and other causes of ARDS. Concomitant FES and pulmonary thromboembolism is a more rarer entity in comparison to FES and pulmonary thromboembolism separately. There are few cases reported in literature wherein patients developed concomitant FES and pulmonary embolism. Almost in all cases fat embolism syndrome and pulmonary embolism occurred either simultaneously or pulmonary thromboembolism was complicated by FES. We report a case wherein patient initially developed FES pre operatively which was later on complicated by acute pulmonary thromboembolism in the post operative period.

Key Words

Fat Embolism syndrome, Pulmonary thromboembolism, Femoral Shaft fracture

Introduction

Fat embolism syndrome is an uncommon but fatal complication usually in orthopedic trauma especially after a long bone fracture. A high level of suspicion should be kept in mind when a patient of long bone fracture develops hypoxia, confusion or rash in the absence of infection and other causes of ARDS. Concomitant FES and pulmonary thromboembolism is a more rarer entity in comparison to FES and pulmonary thromboembolism separately. There are few cases reported in literature wherein patients developed concomitant FES and pulmonary embolism. Almost in all cases fat embolism syndrome and pulmonary embolism occurred either simultaneously or pulmonary thromboembolism was

complicated by FES. We report a case wherein patient initially developed FES pre operatively which was later on complicated by acute pulmonary thromboembolism in the post operative period.

Case Report

A 23 year old young healthy female was brought to orthopedic emergency department with history of road traffic accident. On assessment she had closed bilateral femoral shaft fractures (*Fig 1*). The fractures were immediately splinted and she was admitted for definitive fixation. 48 hours after admission she developed sudden and progressive shortness of breath and hypoxia with an oximetric saturation of 75% on ambient air, requiring *Fio*2

Department of General Medicine, SKIMS Medical College and Hospital, Bemina Srinagar (Jammu and Kashmir) India.

Correspondence to: Dr Aamir Shafi Senior resident, Department of General Medicine, SKIMS Medical College and Hospital, Bemina Srinagar (Jammu and Kashmir) India.

Manuscript Received: 24.06.2022; Revision Accepted: 08.10.2022;

Published Online First: 10 April 2023

Open Access at: <https://journal.jkscience.org>

Copyright: © 2023 JK Science. This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License, which allows others to remix, transform, and build upon the work, and to copy and redistribute the material in any medium or format non-commercially, provided the original author(s) and source are credited and the new creations are distributed under the same license.

Cite this article as: Shafi T, Islam JU, Ahad WA, Zahoor A. Fat Embolism Syndrome Complicated By Acute Pulmonary Thromboembolism After Bilateral femoral shaft Fractures: Two Nightmares In The Same Patient. JK Science 2023;25(2):128-130



Fig. 1 a&b Fracture Bilateral Shaft of Femur and Fracture Fixated with Intramedullary Nail




Fig. 2 CT Image Shows Ground Glass Opacities in bilateral lung fields (R>L)

of 60% to maintain saturation of >94%. There was no altered sensorium, truncal rash, vision abnormality but she was febrile, restless, tachycardic and in respiratory distress. Chest X-ray was grossly normal. Having suffered bilateral femoral shaft fracture and subsequent immobilization a possibility of fat embolism vs pulmonary embolism was made. CTPA was done which showed ground glass opacities in bilateral lung fields more in the lower zones consistent with interstitial

hemorrhage associated with fat embolism (Fig 2). Hemogram revealed drop of hemoglobin from 13 mg/dl to 11 mg/dl and thrombocytopenia. ESR was raised and serum fat macro globules were present fulfilling the GURD and WILSON criteria for Fat Embolism Syndrome. [1] She was managed in ICU with oxygen supR>L)port, aggressive vital monitoring, DVT prophylaxis and intravenous methylprednisolone. She showed progressive improvement in her oxygen saturation over a period of five days and after achieving hemodynamic stability she was subjected to fixation of fractures under spinal anesthesia and immediate post operative course remained uneventful. Three days after internal fixation of fracture she again developed hypoxia and was tachycardic and tachypnoeic on clinical examination. There was no fever or any new infiltrates on chest X-ray. Qualifying as high risk on Well's PTE score CTPA was done which showed thrombi in left interlobar artery and right lower lobe segmental branches (Fig 3a &3b). There was no hypotension or right ventricular strain on echocardiography and cardiac biomarkers were negative. she was started on rivaroxaban 15mg BD. She showed improvement in oxygenation and was discharged after 15 days of complicated hospital course in hemodynamically stable condition on oral




Fig 3a & b Deoicting Thrombus in Left and Thromus in Right Lower Interlobar Artery Segmental

anticoagulation.

Discussion

Fat embolism syndrome and pulmonary embolism are common differential diagnosis in patients of orthopedic trauma and surgeries when they develop hypoxia. [2] However concomitant fat embolism and pulmonary embolism are very rare. After searching extensive literature we found only few cases of concomitant fat and pulmonary embolism. Cothren et al., reported a case of concomitant FES and pulmonary embolism in a patient with pelvic fracture. [3] Randelli *et al.*, reported a case of pulmonary embolism complicated by fat embolism in a patient with bilateral femoral shaft fractures. [4] Shao et al., reported a case of concomitant FES and pulmonary thromboembolism with tympanic membrane perforation in a patient with tibio femoral fracture. [5] Ebinaet al., reported a case of pulmonary thromboembolism complicated by FES in a patient with femoral shaft fracture. [6] Yeak *et al.*, reported a case of fat embolism syndrome and pulmonary embolism in a patient with patent foramen ovale. [7] Our case describes a patient which developed initially fat embolism preoperatively followed by acute pulmonary thromboembolism in the postoperative period.

Conclusion

Concomitant FES and Acute PTE are very rare to occur with few cases reported in literature. We present a case

report in which a patient initially developed FES in pre operative period and later on suffered mild PTE also post operatively. We suggest that patients of orthopedic trauma who develop FES should be closely followed and monitored for the development of PTE. Whether FES is a risk factor for pulmonary thromboembolism needs to be elucidated further in large randomized studies.

References

1. Gurd AR, Wilson RI. The fat embolism syndrome. *J Bone Joint Surg Br* 1974;56B:408-16.
2. Misra I, Glasgow J, Moosavy F. Fat Emboli Syndrome. *Del Med J* 2017 ;89(5):148-150.
3. Cothren CC, Moore EE, Vanderheiden T, Haenel JB, Smith WR. Occam's razor is a double-edged sword: Concomitant pulmonary embolus and fat embolism syndrome. *Journal of Trauma and Acute Care Surgery* 2008 ;65(6):1558-60.
4. Randelli F, Capitani P, Pace F, Favilla S, Galante C, Randelli P. Bilateral femoral shaft fractures complicated by fat and pulmonary embolism: a case report. *Injury* 2015;46:S28-30.
5. Shao J, Kong DC, Zheng XH, Chen TN, Yang TY. Postoperative complications of concomitant fat embolism syndrome, pulmonary embolism and tympanic membrane perforation after tibiofibular fracture: A case report. *World Journal of Clinical Cases.* 2021;9(2):476.
6. Ebina M, Inoue A, Atsumi T, Ariyoshi K. Concomitant fat embolism syndrome and pulmonary embolism in a patient with a femoral shaft fracture. *Acute Medicine & Surgery.* 2016 ;3(2):135-8.
7. Yeak RD, Liew SK. Concomitant fat embolism syndrome and pulmonary embolism in a patient with patent foramen ovale. *Actaorthopaedicaetraumatologicaurcica* 2020 ;54(4):465.